

THE DERMATOPHYTE, *MICROSPORUM GYPSEUM*, AS A SAPROPHYTE AND PARASITE*

LIBERO AJELLO, PH.D.

Vanbreuseghem (1) recently described a selective procedure for isolating keratinophilic fungi from soil. The method extends the work of Karling (2), who first used keratin in various forms as bait for the isolation of chytrids from soil and water. Vanbreuseghem placed hair filaments upon the surface of moistened soil in which *Trichophyton mentagrophytes* (*Ctenomyces interdigitalis*), *T. rubrum* and *Epidermophyton floccosum* had been grown. After a few days, the hair bait became visibly overgrown by mycelium which was noted to be penetrating the hair shafts by means of "perforating organs" (Vanbreuseghem, 3).† Cultures made from the invaded hair yielded *T. rubrum* and *E. floccosum*, but were negative for *T. mentagrophytes*.

Previous attempts to isolate these dermatophytes directly from the test soils had ended in failure, due to overgrowth of the culture tubes by saprophytic molds, which had contaminated the soils during some prior manipulations. Vanbreuseghem examined a number of natural soils with his technic and isolated a keratinophilic fungus which subsequently was designated to be a member of a new genus *Keratinomyces* and given the specific name of *K. ajelloi* (4). However, none of the known dermatophytes was isolated from the Belgian soils studied.

The present paper records the successful isolation of a known dermatophyte from natural soils baited with hair.

MATERIALS AND METHODS

The soils examined in this study were collected from a wide variety of sites in Williamson County, Tennessee, and the counties of Cobb, DeKalb, Gilmer, Rabun and Thomas, Georgia. The Tennessee soils had been used previously in the course of a study of the occurrence and distribution of *Histoplasma capsulatum* in an endemic area of benign histoplasmosis (5-6). All the samples were gathered from the upper surface of the collection site by scooping them directly into 4 oz. bottles.

In the laboratory, sterile Petri dishes were half-filled with the soil specimens, moistened with 15 to 30 ml. of sterile distilled water (the exact amount dependent upon the nature of the soil samples), and baited by placing short tufts of autoclaved human hair upon the surface of the soil. These preparations then were incubated at room temperature (20-25°C) in a dark cupboard and examined periodically for the development of mycelium on the hair filaments.

* From the Communicable Disease Center, U. S. Public Health Service, Department of Health, Education and Welfare, Atlanta, Georgia.

† The radial penetration of hair filaments by the mycelium of dermatophytes was first described by Davidson and Gregory in 1934 (130). These observers clearly noted the formation of pits, extending deep into the hair, as a result of the enzymatic activity of the invading hyphae.

Received for publication April 13, 1953.

Hairs overgrown with mycelium were examined microscopically and cultured on a medium containing cycloheximide* ("acti-dione"), penicillin and streptomycin.

The isolation medium was prepared by adding 0.5 mg. of cycloheximide, 20 units of penicillin and 40 units of streptomycin per ml. of Sabouraud dextrose agar previously sterilized and cooled to 45°C. This medium had been primarily developed for the isolation of *Coccidioides immitis* from soil (7), but was subsequently found to permit the selective isolation of most of the fungi parasitic to man from a variety of heavily contaminated sources (Georg, 8; Ajello and Getz, 9).



FIG. 1. Soil plate showing appearance of *Microsporum gypseum* on hair filaments used as bait. (Soil sample 91A).

RESULTS

Of 74 Tennessee samples examined, 26, or 35.1%, yielded cultures of the dermatophyte *Microsporum gypseum* as did 11 of 42 Georgia specimens (26.2%), giving an over-all recovery of 31.9%.

In all instances, the growth of this dermatophyte upon the bait was vigorous and easily detected with the naked eye (Fig. 1). The hair filaments became covered with a yellowish-white mantle of mycelium. As seen in Fig. 2, the elliptical, echinulate, multiseptate macroconidia, diagnostic of *M. gypseum* were produced in great numbers. In addition, spherical, unicellular microconidia were also produced. The hair filaments were penetrated to varying depths by "perforating organs" (Fig. 3) which were composed of a cone-shaped mass of

* Generously donated by the Upjohn Co., Kalamazoo, Michigan.




FIG. 2. Profuse production of macroconidia by *M. gypseum* on hair filaments exposed to soil (soil isolate G8). Original magnification 100 \times .




FIG. 3. Perforation of hair filament by mycelium of *M. gypseum*. (Isolate G8). Original magnification 475 \times .




FIG. 4. Cultural appearance of *M. gypseum* on Sabouraud dextrose agar (Soil isolate G8)




FIG. 5. Macroconidia of *M. gypseum* as produced on hair (Soil isolate G8). Original magnification 450 \times .




FIG. 6. Appearance of *Keratinomyces ajelloi* on hair filaments used as bait. (Soil sample G18).




FIG. 7. Profuse production of macroconidia by *K. ajelloi* on hair filament. (Soil isolate G39). Original magnification 225X.

mycelium that grew in size as it progressively penetrated the hair filament, eventually perforating the shaft.

Through use of the selective isolation medium, pure cultures of the 37 isolates of *M. gypseum* were obtained. All produced powdery, cinnamon-brown colored




FIG. 8. Penetration of hair filament by *K. ajelloi* (Soil isolate G18). Original magnification 525 \times .




FIG. 9. Colony of *K. ajelloi* on Sabouraud dextrose agar (Soil isolate 100D)

colonies (Fig. 4) and abundant elliptical thin-walled, multiseptate macroconidia that are so typical of this fungus (Fig. 5).

It was not deemed necessary to determine the pathogenicity of these soil isolates, as Gordon (10) had previously verified on a human subject the infective

nature of a culture of *M. gypseum* directly isolated from one of the Tennessee soil samples.

Microscopic examination of the bait soon after exposure to the moistened soil frequently revealed the presence of phycomycetes representative of the orders Chytridiales and Peronosporales. But these rapidly disappeared and never developed to the extent that their growth became visible to the naked eye. A few fungi other than *M. gypseum* developed frank mycelial growth upon this substratum. These were species of *Fusarium*, *Scopulariopsis*, *Penicillium*, *Sepe-donium*, etc. which could be looked upon as adventitious organisms that merely grew over the hair, never producing perforating organs nor bringing about its disintegration.




FIG. 10. Macroconidia of *K. ajelloi* (Soil isolate G18). Original magnification 450X

Others, representing several different fungi, not only produced abundant mycelium but also formed perforating organs. One of these, represented by 3 isolates, one from Tennessee and 2 from Georgia, was not a dermatophyte and proved to be Vanbreuseghem's *K. ajelloi* (Figs. 6-10). (Keratinolysis is thus not strictly a property of the dermatophytes; despite this, the baiting procedure is remarkably selective.) The New World isolates of *K. ajelloi* are morphologically similar to Vanbreuseghem's but differ in their failure to produce a diffusible pigment.

DISCUSSION

M. gypseum was first encountered by Sabouraud in 1894 (11) but at the time was simply referred to as the "Trichophyton du chien." It was first fully studied

and described by Bodin in 1907 (12), who, following a system of classification based principally on clinical criteria, placed it in the now obsolete genus *Achorion* as *A. gypseum*. In subsequent years, this parasite was reisolated by several investigators, who, failing to relate their isolates to Bodin's, gave it new names. The synonymy was lengthened still further by those who made untenable taxonomic improvisations so that today we have the lengthy synonymy presented in Table I. Not until 1928 was Bodin's fungus correctly placed in the genus *Microsporum* by Guiart and Grigorakis (13).

Dermatologists customarily have divided the members of the genus *Microsporum* (*M. audouini*, *M. canis*, *M. gypseum*) into two groups based upon their relative prevalence as parasites of humans and lower animals. Thus, *M. canis*

TABLE I
Synonymy of Microsporum gypseum

<i>Microsporum gypseum</i> (Bodin) Guiart and Grigorakis 1928. (13)
<i>Trichophyton</i> du chien Sabouraud 1894. (11)
<i>Achorion gypseum</i> Bodin 1907. (12)
<i>M. fulvum</i> Sabouraud (Uriburu) 1909. (14, 15)
<i>M. flavescens</i> Horta 1912. (16)
<i>A. serisei</i> Cazalbou 1913. (17)
<i>M. scorteum</i> Priestly 1914. (18)
<i>M. marginatum</i> Cazalbou 1914. (19)
<i>M. xanthodes</i> Fischer 1918. (20)
<i>Sabouraudites gypseus</i> Ota and Langeron 1923. (21)
<i>S. fulvus</i> Ota and Langeron 1923. (21)
<i>S. flavescens</i> Ota and Langeron 1923. (21)
<i>S. xanthodes</i> Ota and Langeron 1923. (21)
<i>Closterospora gypsea</i> Grigorakis 1925. (22)
<i>C. fulva</i> Grigorakis 1925. (22)
<i>Gymnoascus gypseus</i> Nannizzi 1927. (23)
<i>Microsporum</i> sp. Nakamura 1931. (24)
<i>Ectotrichophyton nakamurae</i> Dodge 1935. (25)

and *M. gypseum* are referred to as "animal types" since they have been considered to be primarily parasites of lower animals, such as cats and dogs, which served as sources of human infections. In contrast, *M. audouini* is designated frequently as the "human or anthropophilic type" in the belief that it is predominantly a parasite of man* and infections are transmitted only by contact from person to person.

Of the three *Microsporum* species, *M. gypseum* has long been considered to be the rarest. Reports of *M. gypseum* infections indeed are few, particularly when contrasted with those of *M. audouini* and *M. canis*. A survey of all available records has revealed 155 instances of human infections attributable to *M. gypseum* in the United States and 115 in the remainder of the world (Table II).

* Intensive animal studies may prove otherwise as two instances of canine infections attributed to *M. audouini* are recorded in the literature (Sabouraud, 1908 (26) and Murrell, 1951 (27)).

It is obvious from these data that *M. gypseum* has a global distribution. No basis exists for considering that fungus to be native to Brazil and having been introduced from there into the United States and, inferentially, to other countries as stated by Lewis and Hopper (119).

It was surprising, in view of the alleged association of *M. gypseum* with lower animals, to find recorded only 61 instances of lower animal infections by this fungus (Table III).

The world-wide occurrence of human *M. gypseum* infections, the rarity of cases involving lower animals, and the discovery that this fungus is prevalent in

TABLE II
Geographic Distribution of human Microsporium gypseum infections
(Literature references indicated in parentheses)

NORTH, CENTRAL & SOUTH AMERICA		EUROPE			
Argentina	1 (14)	Austria	9 (90-92)		
Brazil	9 (16, 28-30)	Belgium	1 (93)		
Canada	8 (31-34)	Denmark	20 (94-96)		
Panama	2 (35)	England	14 (97-98)		
Puerto Rico	1 (36)	Finland	3 (99-100)		
United States	155 (37-88)	France	6 (11, 12, 15, 101-103)		
Uruguay	3 (89)	Germany	2 (20, 104)		
Total no. of cases....	179	Hungary	4 (105-106)		
		Italy	4 (107-110)		
		Ireland	1 (111)		
		Netherlands	1 (112)		
		Spain	1 (113)		
		Switzerland	3 (114-115)		
		Total no. of cases...	69		
		AFRICA			
		Belgian Congo	1 (116)	ASIA	
				Japan	1 (24)
AUSTRALIAN AREA					
New Zealand	19 (117-118)	Australia	1 (18)		

World total, 270.

soil leads to the conclusion that soil must be considered the main source of human infection. Lower animals, thus, can no longer be implicated as the prime source of *M. gypseum*. They, like man, are infected from soil. Only infrequently are infections transmitted from animal to animal.

Indications that *M. gypseum* probably existed in soil as a saprophyte had been previously revealed by Mandels, *et al.* (125), who in carrying out laboratory studies on the deterioration of fabrics by molds, isolated this fungus from a piece of wool fabric buried in a soil sample of unknown origin. In 1952, Cooke (126) recovered this dermatophyte from wool buried in 3 soil study plots located

in Idaho and Washington. These observations, although suggestive, did not conclusively prove that *M. gypsum* existed as a saprophyte in soil. However, definitive proof of saprophytism was finally obtained by Gordon, *et al.* (127) when the characteristic macroconidia of *M. gypsum*, which never are produced on tissues of living animals, were demonstrated in a soil sample collected in Tennessee.

TABLE III
Reported animal infections by M. gypsum
(Total 61)

Horse, 50 Belgium, 40 (132) France, 7 (120, 121, 131, 133) Madagascar, 2 (17, 133) German, 1 (122)	Dog, France, (11)
Monkey, 4 South Pacific, 3 (123) <i>Macacus cynomolgus</i> Germany, 1 (124) <i>M. cynomolgus</i>	Cat, 4 Austria, 1 (92) France, 1 (103) Hungary, 1 (106) United States, 1 (70) Tiger, 1 Indochina, 1 (133) Chicken, 1 Italy, 1 (107)

TABLE IV
Correlation of soil habitat with recovery of M. gypsum

HABITAT	NO. OF SAMPLES COLLECTED	NO. POSITIVE FOR <i>M. gypsum</i>	PERCENTAGE POSITIVE
Inside barn.....	9	8	88.8
Barnyard.....	15	10	66.6
Inside chicken house.....	15	2	13.3
Chicken yard.....	7	0	0
Chicken manure.....	4	0	0
Under house.....	14	3	21.4
Near house.....	22	9	40.9
In open.....	13	5	38.5
Bank of stream.....	1	0	0
Forest.....	13	0	0
Other.....	3	0	0
Total.....	116	37	31.9%

An analysis of the sites, from which the soil samples in this study were gathered reveals a significant correlation between the presence of *M. gypsum* in the sample and the collection of the soil from areas frequented by animals (Table IV). Most of the positive collection sites either were obviously populated by animals or were areas frequented by domestic or wild animals and thus presumably seeded with keratinaceous debris.

The observed facts lead one to conclude that *M. gypsum* is essentially a

soil-inhabiting organism. It must be considered as being one of the components of the complex mycoflora of the soil, which in contradistinction to most terricolous molds has the added ability, under appropriate conditions, to invade the keratinaceous tissues of living animals and produce disease. In nature, *M. gypseum* probably plays a highly specialized role, bringing about, along with other keratinophilic organisms, the microbiological breakdown of keratin into simple elements.*

These findings bring to mind the prescient words of Davidson and Gregory (129) who in 1933 stated: "it may well be that cast-off hairs and epidermal scales are the most important natural substrata on which ringworm fungi pass their saprophytic existence and produce spores capable of infecting new human and animal hosts."

SUMMARY

The dermatophyte *Microsporum gypseum* was isolated from 37 of 116 soil samples collected in Tennessee and Georgia.

There was a significant correlation between the positive samples and the presence of animals at the collecting site.

It is suggested that *M. gypseum* is essentially a soil inhabiting fungus that only rarely parasitizes animals.

In nature, *M. gypseum* may be considered to participate in the microbiological breakdown of keratinaceous debris.

Acknowledgment. The writer wishes to express his thanks to Dr. L. D. Zeidberg of the Tennessee Department of Public Health for obtaining certain of the soils used in this study.

REFERENCES

1. VANBREUSEGHEM, R.: Technique biologique pour l'isolement des dermatophytes du sol. Ann. Soc. Belge de méd. trop., **32**: 173-178, 1952.
2. KARLING, J. S.: Keratinophilic chytrids. 1. Rhizophydium keratinophilum n. sp., a saprophyte isolated on human hair and its parasite, Phlyctidium mycetophagum n. sp. Amer. J. Bot., **33**: 751-757, 1946.
3. VANBREUSEGHEM, R.: Lésions déterminées in vitro par les dermatophytes sur des cheveux isolés. C. R. Soc. Biol., **143**: 1302-1303, 1949.
4. ———: Intérêt théorique et pratique d'un nouveau dermatophyte isolé du sol: Keratinomyces ajelloi gen. nov., sp. nov. Bull. Acad. roy. de méd. de Belgique, **38**: 1068-1077, 1952.
5. AJELLO, L. AND ZEIDBERG, L. D.: Isolation of *Histoplasma capsulatum* and *Allescheria boydii* from soil. Science, **113**: 662-663, 1951.
6. ZEIDBERG, L. D., AJELLO, L., DILLON, A. AND RUNYON, L. C.: Isolation of *Histoplasma capsulatum* from soil. Am. J. Pub. Health, **42**: 930-935, 1952.
7. GEORG, L. K., AJELLO, L. AND GORDON, M. A.: A selective medium for the isolation of Coccidioides immitis. Science, **114**: 387-389, 1951.
8. GEORG, L. K.: Use of a cycloheximide medium for the isolation of dermatophytes from clinical materials. Arch. Dermat. & Syph. **67**: 355-361, 1953.

* The keratinolytic power of *M. gypseum* has been demonstrated *in vitro* by Mandels *et al.* (125) and Page (128).

9. AJELLO, L. AND GETZ, M.: Dermatophytes and shoes. J. Invest. Dermat. In press.
10. GORDON, M. A.: The occurrence of the dermatophyte, *Microsporum gypseum* as a saprophyte in soil. J. Invest. Dermat., **20**: 201-206, 1953.
11. SABOURAUD, R.: Les trichophyties humaines. Rueff et Cie. Paris, 1894.
12. BODIN, E.: SuRun nouveau champignon du favus (Achorion gypseum). Ann. de dermat. et syph., **8**: 585-602, 1907.
13. GUIART, J. AND GRIGORAKIS, Z.: La classification botaniques des teignes. Lyon Med., **141**: 369-378, 1928.
14. URIBURU, J. V.: Contribución al estudio de las tiñas en Buenos Aires. Argentina Medica, **7**: 563-582, 1909.
15. SABOURAUD, R.: Les Teignes Masson et Cie., Paris, 1910.
16. HORTA, B.: Contribuições para o estudo das dermatomicozes no Brazil. I. Microsporon flavescentes n. sp. agente duma nova tinha microsporia. Mem. Inst. Oswaldo Cruz., **3**: 301-308, 1912.
17. CAZALBOU, L.: Note sur un nouveau favus du cheval observé à Madagascar. Bull. Soc. path. exot., **6**: 300-303, 1913.
18. PRIESTLY, H.: Microsporon scorteum (n. sp.) from a case of ringworm in man. Ann. Trop. Med., **8**: 113-118, 1914.
19. CAZALBOU, L.: Sur les teignes microsporiques des équides. Deux Microsporon nouveaux. Premier. Congr. Internat. Path. Comp., **2**: 306-308.
20. FISCHER, W.: Sycosis parasitaria durch ein neues Mikrosporum von Tiertype (Microsporum xanthodes). Dermat. Wchnschr., **66**: 241-247, 1918.
21. OTA, M. AND LANGERON, M.: Nouvelle classification des dermatophytes. Ann. de parasitol. Hum. Comp., **1**: 305-336, 1923.
22. GRIGORAKIS, L.: Recherches cytologiques et taxonomiques sur les dermatophytes et quelques autres champignons parasites. Ann. Sci. Nat. Bot., **7**: 165-444, 1925.
23. NANNIZZI, A.: Ricerche sull'origine saprofitica dei funghi delle tigne il Gymnoascus gypseum sp. n., forma ascofora del Sabouraudites (Achorion) gypseum (Bodin). Ota et Langeron. Atti. R. Accad. Fisiocrit., **2**: 89-97, 1927.
24. NAKAMURA, T.: Über eine bisher nicht beschrieben Art von Microsporon in Japan. Jap. Jour. Derm. Urol., **31**: 1462-1472, 1931.
25. DODGE, C. W.: Medical Mycology. C. V. Mosby Co., St. Louis, 1935.
26. SABOURAUD, R.: The dermatomycoses common to man and animals. Brit. M. J., **2**: 1089-1094, 1908.
27. MURRELL, T. W.: Microsporum audouini. Isolated from a dog. Arch. Dermat. & Syph., **63**: 638, 1951.
28. MARTINS DE CASTRO, A.: Achorion gypseum, Bodin 1907. Am. Bras. Derm. Sif., **14**: 1-14, 1939.
29. ROSSETTI, N.: Contribuição para estudo do Achorion gypseum Bodin, 1907. Quatro casos observados em São Paulo. Arq. de dermat. e sif. de São Paulo, **3**: 3-45, 1939.
30. DEALMEIDA, F., LACAZ, C. S. AND COSTA, O.: Dados estatísticos sobre as principais micoses humanas observados em nosso meio. Annais Fac. Med. Univ. São Paulo, **24**: 39-62, 1948-49.
31. BURGESS, J. W.: Fungus infections of the skin. Report on cultural study of the flora of ringworm. Arch. Dermat. & Syph., **12**: 835-857, 1925.
32. DANBY, C. W. E. AND FORSEY, R.: A cultural study of ringworm. Canad. M. A. J., **56**: 640-642, 1947.
33. FISHER, J. B. AND WRONG, N. M.: Fungous infections of the skin, hair and nails. Canad. M. A. J., **67**: 398-403, 1952.
34. BIRT, A. R.: Personal communication, 1952.
35. SHRAGER, J.: Personal communication, 1952.
36. CARRION, A. L. AND SILVA, M.: Ringworm of the scalp in Puerto Rico. Puerto Rico J. Pub. Health & Trop. Med., **20**: 320-391, 1944.
37. MEWBORN, A. D.: Report of a case of favus of scrotum, coexisting with ringworm of the thigh, giving identical trichophyton-like cultures. Jour. Cut. Dis., **21**: 11-18, 1903.

38. SNOW, J. S. AND LONDON, I. D.: Management of ringworm of the scalp. Present status of the epidemic in the Birmingham area. *J. M. A. Alabama*, **17**: 333-336, 1948.
39. ULRICH, E. W.: Personal communication, 1952 (Arkansas).
40. WILSON, J. W. AND PUNKETT, O. A.: Lack of fluorescence of scalp hairs infected with *Microsporum gypseum* (fulvum). *J. Invest. Dermat.*, **16**: 119-120, 1951.
41. BROWNE, A. S.: Personal communication. 1952. (California).
42. FREEMAN, E.: Personal communication. 1952. (Florida).
43. DOBES, W. L.: Treatment of tinea capitis with estrogenic hormones. *Arch. Dermat. & Syph.*, **62**: 58-65, 1950.
44. MORRIS, J.: Personal communication. 1952. (Georgia).
45. AJELLO, L.: Unpublished data (Georgia, Louisiana, Maryland, Ohio).
46. BENEDEK, T. AND FELSHER, I. M.: Epidemiology of tinea capitis. I. A study of tinea capitis in a dispensary. *Arch. Dermat. & Syph.*, **49**: 120-123, 1944.
47. DALTON, J. E., SLAUGHTER, J. C., JENKINS, R. E., PHELPS, S. AND HACKNEY, V. C.: Microsporiasis due to *Microsporum gypseum*. *J. Invest. Dermat.*, **15**: 421-423, 1950.
48. DAMON, S. R.: Personal communication. 1952. (Indiana).
49. CARNEY, R. G.: Inflammatory ringworm due to *Trichophyton faviforme*. Report of seventeen cases. *Arch. Dermat. & Syph.*, **59**: 209-216, 1949.
50. HARRIS, H. L.: Personal communication. 1952. (Kansas).
51. SUTTON, R. L., JR.: Personal communication. 1952. (Kansas).
52. DALTON, J. E., SLAUGHTER, J. C., JENKINS, R. E., PHELPS, S. AND HACKNEY, V. C.: Microsporiasis due to *Microsporum gypseum*. *J. Invest. Dermat.*, **15**: 421-423, 1950.
53. GALTHER, G. M. AND FRIESHABER, F. C.: Tinea capitis. A survey. *New Orleans M. & S. J.*, **103**: 256-258, 1950.
54. SCHWARTZ, L., PECK, S. M., BOTVINICK, I., LEIBOVITZ, A. L. AND FRASIER, E. S.: Control of ringworm of scalp among school children in Hagerstown, Maryland. 1944-45. *Pub. Health Bull.*, **294**: 1-27, 1946.
55. CONANT, N. F.: Studies in the genus *Microsporum* III Taxonomic studies. *Arch. Dermat. & Syph.*, **36**: 781-808, 1937.
56. SWARTZ, J. H., ROCKWOOD, E. M. AND GLICKLIGH, E. A.: A survey of tinea capitis including favus. *Arch. Dermat. & Syph.*, **60**: 486-494, 1949.
57. DOWNING, J. G., BAIRD, J. W. AND PACI, D.: Fungus diseases in the Boston area. *New England J. Med.*, **243**: 594, 1950.
58. CAWLEY, E. P. AND GREKIN, R. H.: Parafavus restricted to the scrotum. Report of a case. *Arch. Dermat. & Syph.*, **60**: 435-436, 1949.
59. FISHER, I.: Personal communication. 1952. (Minnesota).
60. RAVITS, H. G.: Personal communication. 1952. (Minnesota).
61. WILLIAMS, F. S.: Personal communication. 1952. (Mississippi).
62. MOORE, M. AND CONRAD, A. H.: Microsporiasis of the scalp caused by *Microsporum fulvum*. *Arch. Dermat. & Syph.*, **42**: 610-613, 1940.
63. SUTTON, R. L., JR.: Personal communication. 1952. (Missouri).
64. LOBITZ, W. C.: Personal communication. 1952. (New Hampshire).
65. EMMONS, C. W.: Observations on *Achorion gypseum*. *Mycologia*, **23**: 87-95, 1931.
66. LEWIS, G. M. AND HOPPER, M. E.: An introduction to medical mycology. The Yearbook Publishers, Inc., Chicago, 1948.
67. MONTGOMERY, R. M., HEINLEIN, J. A. AND KARPLUK, F. E.: Ringworm of the scalp in New York. *N. Y. State Jour. Med.*, **48**: 629-630, 1949.
68. CULBERT, R. W., ROBINSON, A. E. R. AND LERNER, M. N.: Study in the reduction of absence from school of children with tinea capitis. *Am. J. Pub. Health*, **40**: 1089-1095, 1950.
69. BENHAM, R.: Personal communication. 1952. (New York).
70. LAMB, J.: Personal communication. 1952. (Oklahoma).
71. LIVINGOOD, C. S. AND PILLSBURY, D. M.: Ringworm of the Scalp: Prolonged observations, family investigation, cultural and immunologic studies in 130 cases. *J. Invest. Dermat.*, **4**: 43-57, 1941.

72. KLIGMAN, A. M.: Personal communication. 1952. (Pennsylvania).
73. SCULLY, J. P.: Personal communication. 1952. (Pennsylvania).
74. ALLISON, A. M.: Personal communication. 1952. (South Carolina).
75. ADAMS, A. S.: Personal communications. 1952. (South Carolina).
76. BUCHANAN, R. N.: Personal communication. 1952. (Tennessee).
77. HALL, E. R.: Personal communication. 1952. (Tennessee).
78. MILLER, F.: Personal communication. 1952. (Tennessee).
79. LEHMAN, C. F., PIPKIN, J. L. AND ROSSMAN, A. C.: Cultural survey of tinea capitis in San Antonio, Texas. Arch. Dermat. & Syph., **61**: 488, 1950.
80. SHARP, W. B. AND WEGNER, M. J.: Microsporum gypseum as an etiologic agent of tinea in the United States. Arch. Derm. & Syph., **61**: 824-826, 1950.
81. SEALE, E.: Personal communication. 1952. (Texas).
82. FLOWER, A. H., JR.: Personal communication. (Vermont).
83. SMITH, G. C.: Personal communication. 1952. (Virginia).
84. TRICE, E. R. AND SHAFER, J. C.: Occurrence of Microsporum gypseum (*M. fulvum*) infections in the District of Columbia area. Arch. Dermat. & Syph., **64**: 309-313, 1951.
85. HATHAWAY, J. C.: Personal communication. 1952. (Washington).
86. POTTER, R. T.: Personal communication. 1952. (Washington).
87. JOHNSON, S. A. M.: Personal communication. 1952. (Wisconsin).
88. RUCH, D. M.: Personal communication. 1952. (Wisconsin).
89. MACKINNON, J. E.: Estadística sobre 1000 casos de micosis cutáneas en el Uruguay y determinación de las especies causantes. An. Inst. Hig. Montevideo, **3**: 83-94, 1949.
90. STEIN, R. O.: Über cutireaktion bei favus. Arch. Dermat. & Syph., **132**: 294-303, 1921.
91. TZSCHIRNTSCH, K.: Über favus corporis durch Achorion gypseum. Derm. Zeitschr., **51**: 331-335, 1927.
92. SCHERBER: Verhandlungen der Wiener dermatologische Gesellschaft. Arch. f. Derm. u. Syph., **125**: 47, 1920.
93. LEFEVRE, A.: Cited by Sabouraud, R. 1910. Les Teignes. (pg. 573), Masson et Cie, Paris, 1908.
94. RASCH, C.: Skin affections caused by Achorion gypseum (Bodin). Brit. J. Dermat. & Syph., **34**: 1-5, 1922.
95. SYLVEST, B.: The incidence of dermatophytes in Denmark. Acta. dermat.-venereol., **29**: 225-232, 1949.
96. HAXTHAUSON, H.: Treatment of superficial trichophytosis with CO₂-snow. An attempt at artificial immunisation ("imitated kerion"). Acta. dermat.-venereol., **30**: 405-416, 1950.
97. WALKER, J.: The dermatophytoses of Great Britain. Brit. J. Dermat. & Syph., **62**: 239-251, 1950.
98. CALNAN, C. D.: Ringworm of the scalp. Brit. J. Dermat., **64**: 343-344, 1952.
99. HÄRÖ, S.: On the asymptomatic occurrence of ringworm of the feet in Finland. Ann. Med. Exp. Biol. Fenn., **27**: 73-77, 1949.
100. PÄTILÄ, R. AND HÄRÖ, S.: Review of fungi found on the skin on the basis of the 1948 material. Karestenia, **1**: 48-59, 1950.
101. SABRAZES, J. AND BRENGUES.: Trichophytie profonde de la barbe. Etude historique, clinique et anatomopathologique. Actes Soc. Linn. Bordeaux, **53**: 211-220, 1898.
102. RABUT, R. ET DUMET: Un cas de "favus gypseum" de Bodin. Bull. Soc. Franc. Derm. Syph., **30**: 391-393, 1923.
103. RABUT, R. ET CANAL: Deux foyers de Microsporum felenium. Un cas d'Achorion gypseum. Bull. Soc. Franc. Derm. Syph., **34**: 891-892, 1927.
104. MÜLLER, H.: Versammlung südwestdeutscher Dermatologen. Zbl. f. Haut. u. Geschl., **13**: 28-31, 1924.
105. NEUBER, E.: Die in Ungarn vorkommenden pathogenen Pilze auf der Haut und ihren anhängen. Eine neue Trichophytieart, Trichophyton fuscum sulcatum. Derm. Woch., **80**: 861-871, 1925.

106. BALLAGI, S.: Die in Ungarn einheimischer Mikrosporon, Trichophyton-, Epidermophyton-, and Achorion-pilze. *Derm. Woch.*, **83**: 1155-1169, 1926.
107. TRUFFI, M.: Favo della palpebra da Achorion gypseum. *Giorn. Ital. Mal. Ven. Pelle.*, **55**: 330-331, 1914.
108. TRUFFI, M.: Favo da achorion gypseum. *Giorn. Ital. Ven. Pelle.*, **64**: 557-563, 1923.
109. MAZZINI, L.: Un caso di favo de Achorion gypseum (Bodin). *Arch. Ital. Derm. Sifiligr. Venereal.*, **1**: 277-280, 1925.
110. CARLI, G.: Favo epidermico dello scroto ed eritemato squamoso della cute glabra. *Arch. Ital. Derm.*, **22**: 49-53, 1949.
111. MEENAN, F. O. C.: Ringworm species in Eire. *Irish Jour. Med. Sci.*, **311**: 546-549, 1951.
112. BEINTEMA, K.: Microsporon xanthodes or Achorion gypseum. *Nederl. tijdschr. v. genesk.*, **2**: 3326, 1929.
113. PEREIRO MIGUENS, M.: Personal communication, 1952.
114. BORNHAUSER, S.: Über einige relativ seltene hautpathogene Pilze. *Dermatologica*, **101**: 345-356, 1950.
115. BLANK, F.: Zur Dermatophytien Flora der Schwiez. *Dermatologica*, **102**: 88-91, 1951.
116. VANBREUSEGHEM, R. AND BORGERS, G.: A propos d'une souche de Sabouraudites (Microsporum) gypseus isolée au Congo belge. *Ann. Soc. Belge Med. Trop.*, **21**: 377-382, 1951.
117. MARPLES, M. J.: The diagnosis of fungus infection. An investigation of 100 cases. *New Zealand Med. Jour.*, **46**: 422-427, 1947.
118. MARPLES, M. J.: Some observations on the occurrence and clinical course of tinea capitis and corporis in Otago. *New Zealand M. J.*, **50**: 460-479, 1951.
119. LEWIS, G. M. AND HOPPER, M. E.: An Introduction to Medical Mycology. The Year-book Pub., Inc. Chicago, 1948.
120. SUIJS, A.: Cited by Sabouraud, R. 1910. Les Teignes (pg. 573) Mason et Cie., Paris, 1908.
121. URBAIN, A., BAROTTO, J. AND CAPDEBEILLE, H. B.: Sur un cas de teigne equine due a l'achorion gypseum. *Bull. Mem. Soc. Centr. Med. Veterinaire*, **102**: 50, 1926.
122. BERGNER, K.: Mikrosporon equinum und Achorion gypseum als eneger von Flechten-erkrankungen bei Pferden. *Z. Infekkr. Haustiere*, **58**: 121-141, 1942.
123. HASEGAWA, M. AND YAMAMOTO, K.: Über einen bei Affen gezuchteten Pilz: Microsporon fulvum. *Jap. J. Dermat. & Urol.*, **39**: 220-222, 1936 (In Japanese).
124. HOFFMAN, H.: Über eine Pilzerkrankung beim Affen hervorgerufen wahrscheinlich durch Achorion gypseum. *Derm. Woch.*, **86**: 353-354, 1928.
125. MANDELS, G. R., STAHL, W. H. AND LEVINSON, H. S.: Structural changes in wool degraded by the ringworm fungus Microsporum gypseum and other microorganisms. *Textile Rec. Jour.*, **18**: 224-231, 1948.
126. COOKE, W. B.: Western Fungi—II. *Mycologia*, **44**: 245-261, 1952.
127. GORDON, M. A., AJELLO, L., GEORG, L. K. AND ZEIDBERG, L. D.: Microsporum gypseum and Histoplasma capsulatum spores in soil and water. *Science*, **116**: 208, 1952.
128. PAGE, R. M.: Observations on keratin digestion by Microsporum gypseum. *Mycologia*, **42**: 591-602, 1950.
129. DAVIDSON, A. M. AND GREGORY, P. H.: Development of fuseaux, aleuriospores and spirals on detached hairs infected by ringworm fungi. *Nature*, **131**: 836-837, 1933.
130. ———: In situ cultures of dermatophytes. *Can. Jour. Res.*, **10**: 373-393, 1934.
131. BROCC-ROUSSEU, D., URBAIN, A. AND BAROTTE, J.: Etude des teignes du cheval et l'immunité dans les teignes experimentales. *Ans. Inst. Pasteur* **41**: 513-551, 1927.
132. NEEFS AND GILLAIN: Contribution à l'étude de la teigne. *Ann. Med. Vet. Belge* **76**: 193-209, 1931.
133. LEBASQUE, J.: Les champignons des teignes du cheval et des bovides. Thèse. Librairie Gouillet, Paris, 1933.